

On Long Range Percolation With Heavy Tails

S. Friedli¹, B.N.B. de Lima², V. Sidoravicius³

Abstract

Consider independent long range percolation on \mathbf{Z}^2 , where horizontal and vertical edges of length n are open with probability p_n . We show that if $\limsup_{n \rightarrow \infty} p_n > 0$, then there exists an integer N such that $P_N(0 \leftrightarrow \infty) > 0$, where P_N is the truncated measure obtained by taking $p_{N,n} = p_n$ for $n \leq N$ and $p_{N,n} = 0$ for all $n > N$.

Keywords and phrases: Long range percolation, truncation, slab percolation.
Mathematics Subject Classification (2000): Primary 60K35, 82B44.

On the graph $\mathcal{G} = (\mathcal{V}, \mathcal{E})$ with $\mathcal{V} = \mathbf{Z}^2$, and $\mathcal{E} = \{\langle x, y \rangle \subset \mathcal{V} \times \mathcal{V} : |x_1 - y_1| = 0, \text{ or } |x_2 - y_2| = 0\}$, consider a long range percolation process (Ω, \mathcal{F}, P) , where $\Omega = \{0, 1\}^{\mathcal{E}}$, $P = \prod_{\langle x, y \rangle \in \mathcal{E}} \mu_{\langle x, y \rangle}$, and $\mu_{\langle x, y \rangle} \{\omega_{\langle x, y \rangle} = 1\} = p_{|x-y|} \in [0, 1]$ is a Bernoulli measure, independent of the state of other edges. Given a sequence $(p_n)_{n \in \mathbf{N}}$ and $N \in \mathbf{N}$, we define a truncated sequence $(p_{N,n})_{n \in \mathbf{N}}$ by

$$p_{N,n} = \begin{cases} p_n & \text{if } n \leq N, \\ 0 & \text{if } n > N, \end{cases} \quad (1)$$

and a truncated percolation process by taking $P_N = \prod_{\langle x, y \rangle \in \mathcal{E}} \mu_{N, \langle x, y \rangle}$, where $\mu_{N, \langle x, y \rangle} \{\omega_{\langle x, y \rangle} = 1\} = p_{N, |x-y|}$.

In this note we adress the following question: given a sequence $(p_n)_{n \in \mathbf{N}}$ for which $P(0 \leftrightarrow \infty) > 0$, does there always exists some large enough N such that $P_N(0 \leftrightarrow \infty) > 0$? In other words, given a system with infinite range translation invariant interactions which exhibits a phase transition, we ask if the infiniteness of the range is really crucial for this transition to occur. It is known, for instance, that infinite range is essential in one dimensional systems (cf. [FS], [NS]), but it is believed that in dimensions $d \geq 2$, occurrence (if so) of phase transitions for translation invariant interactions is *always* determined by a bounded part of the interaction (excluding cases when interactions are of intrinsically one dimensional structure). Returning to the percolation case, rapid (say, exponential) decay or summability of the p_n 's indicates that long range connections may not be necessary for the existence of an infinite cluster. This is the setup of [MS] and partially [B]. On the other hand, heavy tail

¹CBPF, Rua Dr. Xavier Sigaud, 150, Urca, CEP 22290-180, Rio de Janeiro, and IMPA, Estrada Dona Castorina, 110 Jardim Botânico CEP 22460-320, Rio de Janeiro.

²UFMG, Av. Antônio Carlos 6627 C.P. 702 Belo Horizonte 30123-970.

³IMPA, Estrada Dona Castorina, 110 Jardim Botânico CEP 22460-320, Rio de Janeiro.

interactions are still poorly understood, and the only existing studies, [SSV] and [B], rely heavily on asymptotic monotonicity assumptions which are a key ingredient for the use of rather laborious coarse-graining techniques. Although the approach we present here relies on deep and highly nontrivial facts ([GM], [K]), it leads to a much shorter (not to say elementary) proof, which is less sensitive to the geometry of the interactions, and allows to consider a rather general class of systems with connections of irregular, in particular lacunary structure.

Theorem 1. ($d \geq 2$) *If $\limsup_{n \rightarrow \infty} p_n > 0$, then*

$$P_N(0 \leftrightarrow \infty) > 0 \quad (2)$$

for some large enough N .

Proof. It suffices to consider $d = 2$. Define $\epsilon > 0$ by $2\epsilon = \limsup_{n \rightarrow \infty} p_n > 0$. By [K], Theorem 1 p. 220, there exists d_ϵ such that

$$p_c(\mathbf{Z}^{d_\epsilon}) < \epsilon/2, \quad (3)$$

and by [GM], Theorem A p. 447, we can find K_ϵ such that

$$p_c(\{1, 2, \dots, K_\epsilon\}^{d_\epsilon-2} \times \mathbf{Z}^2) < p_c(\mathbf{Z}^{d_\epsilon}) + \epsilon/2 < \epsilon. \quad (4)$$

Let $n_0 = 0$. For $j \in \{1, 2, \dots, d_\epsilon - 1\}$, define recursively

$$n_j = \min\{\ell > (K_\epsilon + 1)n_{j-1} : p_\ell \geq \epsilon\}.$$

For $x \in \mathbf{Z}^2$ and $B \subseteq \mathbf{Z}^2$, define $\mathbf{T}_x B = \{z + x, z \in B\}$. Set $B_0 = \{(0, 0)\}$. For $j \in \{1, 2, \dots, d_\epsilon - 2\}$, define $B_j = \cup_{m=0}^{K_\epsilon-1} \mathbf{T}_{m(n_j, 0)} B_{j-1}$. Then, let

$$\mathcal{V}_{d_\epsilon-1} = \bigcup_{(k, m) \in \mathbf{Z}^2} \mathbf{T}_{(kn_{d_\epsilon-1}, mn_1)} B_{d_\epsilon-2},$$

and

$$\begin{aligned} \mathcal{E}_{d_\epsilon-1} = \{ \langle x, y \rangle, x, y \in \mathcal{V}_{d_\epsilon-1} : |x_1 - y_1| = n_j \text{ for some } 1 \leq j \leq d_\epsilon - 1 \\ \text{and } x_2 = y_2, \text{ or } x_1 = y_1 \text{ and } |x_2 - y_2| = n_1 \}. \end{aligned}$$

It is straightforward that the graph $\mathcal{G}_{d_\epsilon-1} = (\mathcal{V}_{d_\epsilon-1}, \mathcal{E}_{d_\epsilon-1})$ is isomorphic to $\{1, 2, \dots, K_\epsilon\}^{d_\epsilon-2} \times \mathbf{Z}^2$. Moreover, by our choice of n_j , $1 \leq j \leq d_\epsilon - 1$, we have that each edge of $\mathcal{G}_{d_\epsilon-1}$ is open with probability at least ϵ , and using (4) we get (2) with $N = n_{d_\epsilon-1}$. \square

Remark. For further applications of this method to percolation and interacting spin systems see [FL].

Acknowledgments. S.F. is supported by the Fonds National Suisse pour la Recherche Scientifique, B.N.B.L. and V.S. are partially supported by CNPq and FAPERJ. F.S. wishes to thank CBPF and IMPA for hospitality and support, B.N.B.L. wishes to thank IMPA for hospitality and financial support during multiple visits.

References

- [B] Berger N., *Transience, Recurrence and Critical Behavior for Long-Range Percolation*, Commun. Math. Phys. **226**, 531-558 (2002).
- [FL] Friedli S., de Lima B.N.B., in preparation.
- [FS] Fröhlich J., Spencer B. *The phase transition in the one-dimensional Ising model with $1/r^2$ interaction energy*, Comm. Math. Phys. **84**, no. 1, 87–101 (1982).
- [GM] Grimmett G., Marstrand J.M., *The Supercritical Phase of Percolation is Well Behaved*, Proc. Roy. Soc. London Ser **A 430**, 439-457 (1990).
- [K] Kesten H., *Asymptotics in high dimensions for percolation*, Disorder in physical systems, 219–240, Oxford Sci. Publ., Oxford Univ. Press, New York, 1990.
- [MS] Meester R., Steif J.E., *On the Continuity of the Critical Value for Long Range Percolation in the Exponential Case*, Commun. Math. Phys. **180**, 483-504 (1996).
- [NS] Newman C.M., Schulman L.S., *One Dimensional $1/|i - j|^s$ Percolation Models: The Existence of a Transition for $s \leq 2$* , Commun. Math. Phys. **104**, 547-571 (1986).
- [SSV] Sidoravicius V., Surgailis D., Vares M.E., *On the Truncated Anisotropic Long-Range Percolation on \mathbf{Z}^2* , Stoch. Proc. and Appl. **81**, 337-349 (1999).